

Representations of real numbers induced by probability distributions on \mathbb{N}

Jörg Neunhäuserer
Leibnitz University of Hannover, Germany
joerg.neunhaeuserer@web.de

Abstract

We observe that a probability distribution supported by \mathbb{N} , induces a representation of real numbers in $[0, 1)$ with digits in \mathbb{N} . We first study the Hausdorff dimension of sets with prescribed digits with respect to these representations. Then we determine the prevalent frequency of digits and the Hausdorff dimension of sets with prescribed frequencies of digits. As examples we consider the geometric distribution, the Poisson distribution and the zeta distribution.

MSC 2010: 11K55, 60E05, 28A78, 28A80

Key-words: representations of real numbers, discrete distributions, digits, frequency, Hausdorff dimension

1 The Representations

Let $\mathbf{p} = (p_i)_{i \in \mathbb{N}}$ be a probability distribution supported by \mathbb{N} , this means $p_i \in (0, 1)$ for all $i \in \mathbb{N}$ and

$$\sum_{n=1}^{\infty} p_n = 1.$$

Let $\widehat{p}_1 = 0$ and

$$\widehat{p}_n = \sum_{i=1}^{n-1} p_i$$

if $n \geq 2$. For $n \in \mathbb{N}$ we consider linear contractions $T_n : [0, 1) \rightarrow [0, 1)$, given by

$$T_n x = p_n x + \widehat{p}_n,$$

and introduce a map $\pi_{\mathbf{p}} : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{R}$ by

$$\begin{aligned} \pi_{\mathbf{p}}((n_j)) &= \lim_{j \rightarrow \infty} T_{n_1} \circ T_{n_2} \circ \cdots \circ T_{n_j}(0) \\ &= \widehat{p}_{n_1} + \sum_{j=1}^{\infty} p_{n_1} \cdots p_{n_j} \widehat{p}_{n_{j+1}}. \end{aligned}$$

The limit in this expressions existence since the maps T_n are contractions, moreover it is easy to see that:

Proposition 1.1 For all probability distributions \mathbf{p} supported by \mathbb{N} , the map $\pi_{\mathbf{p}} : \mathbb{N}^{\mathbb{N}} \rightarrow [0, 1)$ is a bijection. Moreover the map $\pi_{\mathbf{p}}$ is continuous, if we endorse $\mathbb{N}^{\mathbb{N}}$ with the metric

$$d((n_i), (m_i)) = \sum_{i=1}^{\infty} \delta(n_i, m_i) 2^{-i},$$

where $\delta(0) = 0$ and $\delta(x) = 1$ if $x \neq 0$.

Proof. If $n_1 < n_2$ we have $p_{n_1} + \hat{p}_{n_1} \leq p_{n_2}$ and hence

$$T_{n_1}([0, 1)) \cap T_{n_2}([0, 1)) = [\hat{p}_{n_1}, p_{n_1} + \hat{p}_{n_1}) \cap [\hat{p}_{n_2}, p_{n_2} + \hat{p}_{n_2}) = \emptyset.$$

It follows that $\pi_{\mathbf{p}}$ is injective. Moreover

$$\bigcup_{n=1}^{\infty} T_n([0, 1)) = [0, p_1) \cup \bigcup_{n=1}^{\infty} [\sum_{i=1}^n p_i, \sum_{i=1}^{n+1} p_i) = [0, 1)$$

hence $\pi_{\mathbf{p}}$ is surjective. If $d((n_i), (m_i)) < 2^{-u}$ we have $n_i = m_i$ for $i = 1, \dots, u$, which implies

$$|\pi_{\mathbf{p}}((n_i)) - \pi_{\mathbf{p}}((m_i))| < \max\{p_i | i \in \mathbb{N}\}^u.$$

This proves that $\pi_{\mathbf{p}}$ is continuous with respect to d . □

For $x \in [0, 1)$ we call the sequence $\pi_{\mathbf{p}}^{-1}(x)$ in $\mathbb{N}^{\mathbb{N}}$ the representation of x with respect to the probability distribution \mathbf{p} . The entry of this sequence are the digits of x with respect to the representation, given by \mathbf{p} .

Let us look at three examples. For the geometric distribution on \mathbb{N} , given by $p_i = (1 - p)p^{i-1}$ with $p \in (0, 1)$, we obtain

$$\pi_{\mathbf{p}}((n_j)) = (1 - p^{n_1-1}) + \sum_{j=1}^{\infty} (1 - p)^j p^{n_1 + \dots + p_{n_j}} (1 - p^{n_{j+1}-1}).$$

For the Poisson distribution on \mathbb{N} given by $p_i = e^{-\lambda} \lambda^{i-1} / (i-1)!$ with $\lambda > 0$, we have

$$\pi_{\mathbf{p}}((n_j)) = e^{-\lambda} \sum_{i=1}^{n_1-1} \frac{\lambda^{(i-1)}}{(i-1)!} + \sum_{j=1}^{\infty} e^{-\lambda(j+1)} \frac{p^{n_1 + \dots + n_j}}{n_1! \dots n_j!} \sum_{i=1}^{n_{j+1}-1} \frac{\lambda^{(i-1)}}{(i-1)!}.$$

For the zeta distribution, given by $p_i = i^{-s} / \zeta(s)$ on \mathbb{N} with $s > 1$, we find

$$\pi_{\mathbf{p}}((n_j)) = \zeta(s)^{-1} \sum_{i=1}^{n_1-1} i^{-s} + \sum_{j=1}^{\infty} \zeta(s)^{-(j+1)} (n_1 \dots n_j)^{-s} \sum_{i=1}^{n_{j+1}-1} i^{-s}.$$

As far as we know these representations of real numbers were not considered jet.

2 Prescribed digits

Let $D \subseteq \mathbb{N}$ a set of digits. We are interested in the set $\pi_{\mathbf{p}}(D^{\mathbb{N}})$ of real numbers, which have only digits in D in their representation with respect to a probability distribution \mathbf{p} on \mathbb{N} . It turns out that these sets have Lebesgue measure zero if $D \neq \mathbb{N}$. Thus we study the Hausdorff dimension of these sets. Let us recall that the d -dimensional Hausdorff measure of a set $B \subseteq \mathbb{R}$ is given by

$$\mathfrak{H}^d(B) = \liminf_{\epsilon \rightarrow 0} \left\{ \sum_{i=1}^{\infty} (b_i - a_i)^d \mid B \subset \bigcup_{i=1}^{\infty} [a_i, b_i]; \forall i \in \mathbb{N} : (b_i - a_i) \leq \epsilon \right\}$$

and the Hausdorff dimension of B is

$$\dim_H B = \inf\{d \mid H^d(B) = 0\} = \sup\{d \mid H^d(B) = \infty\}.$$

We recommend [1] or [8] as an introduction to dimension theory. Using the notion of Hausdorff dimension, we obtain:

Theorem 2.1 *Let $\mathbf{p} = (p_i)_{i \in \mathbb{N}}$ be a probability distribution supported by \mathbb{N} and $D \subseteq \mathbb{N}$. If $d \geq 0$ is the solution of*

$$\sum_{i \in D} p_i^d = 1,$$

we have $\dim_H \pi_{\mathbf{p}}(D^{\mathbb{N}}) = d$.

Proof. We have

$$\begin{aligned} \bigcup_{i \in D} T_i(\pi_{\mathbf{p}}(D^{\mathbb{N}})) &= \bigcup_{i \in D} T_i \left\{ \lim_{j \rightarrow \infty} T_{n_1} \circ T_{n_2} \circ \cdots \circ T_{n_j}(0) \mid n_j \in A \forall j \in \mathbb{N} \right\} \\ &= \bigcup_{i \in D} \left\{ \lim_{j \rightarrow \infty} T_i \circ T_{n_1} \circ T_{n_2} \circ \cdots \circ T_{n_j}(0) \mid n_j \in A \forall j \in \mathbb{N} \right\} = \pi_{\mathbf{p}}(D^{\mathbb{N}}). \end{aligned}$$

This means that $\pi_{\mathbf{p}}(D^{\mathbb{N}})$ is the attractor of the linear iterated function system $\{T_i \mid i \in A\}$ on $[0, 1)$, see [3] for finite sets D and [2] for infinite sets. The system fulfills the strong open set condition $T_i((0, 1)) \cap T_j((0, 1)) = \emptyset$ for $i \neq j$. If A is finite, the result directly follows from the classical work of Moran [6]. If A is infinite it follows from theory of infinite iterated function systems see theorem 3.11 of [2] or [7] for a more general approach. \square

As a corollary, we obtain an analogon of Jarnik's [4] classical result on continued fractions.

Corollary 2.1 *If B is the set of bounded sequences in $\mathbb{N}^{\mathbb{N}}$, we have $\dim \pi_{\mathbf{p}}(B) = 1$ for all \mathbf{p} .*

Proof. Since Hausdorff dimension is countable stable

$$\dim_H B = \dim_H \bigcup_{k=1}^{\infty} \pi_{\mathbf{p}}(\{1, \dots, k\}^{\mathbb{N}}) = \sup\{\dim_H \pi_{\mathbf{p}}(\{1, \dots, k\}^{\mathbb{N}}) | k \in \mathbb{N}\} = 1.$$

□

In the following we consider $D = \{1, \dots, n\}$ and $\Pi_{\mathbf{p}}(n) = \pi_{\mathbf{p}}(\{1, \dots, n\}^{\mathbb{N}})$. If \mathbf{p} is the geometric distribution with $p \in (0, 1)$, we obtain $\dim_H \Pi_{\mathbf{p}}(n) = d$, where d is the solution of

$$(1 - p^{dn})(1 - p)^d / (1 - p^d) = 1.$$

We list the first digits of d for some n and p in the following table:

n/p	0.1	0.25	0.5	0.75	0.9
2	0.96875	0.88920	0.69424	0.45439	0.29434
3	0.99718	0.97718	0.87914	0.66352	0.45656
4	0.99972	0.99463	0.94677	0.77979	0.56428
5	0.99997	0.99868	0.97522	0.85084	0.64218
6	0.99999	0.99967	0.98810	0.89611	0.70137

For all $n \geq 2$ we have

$$\lim_{p \rightarrow 0} (1 - p^{1 \cdot n})(1 - p)^1 / (1 - p^1) = 1 \text{ and } \lim_{p \rightarrow 1} (1 - p^{dn})(1 - p)^d / (1 - p^d) = 0 \quad \forall d > 0,$$

hence the dimension attains all values in $(0, 1)$ for $p \in (0, 1)$ by continuity.

Now let \mathbf{p} be the Poisson distribution with $\lambda > 0$. We have $\dim_H \Pi_{\mathbf{p}}(n) = d$, where d is the solution of

$$\sum_{i=1}^n e^{-\lambda d} \lambda^{d(i-1)} / ((i-1)!)^d = 1.$$

We again list the first digits of d for some n and λ in a table:

n/λ	0.25	0.5	1	2	4
2	0.94980	0.87189	0.69314	0.42577	0.21288
3	0.99642	0.98345	0.92666	0.73178	0.40665
4	0.99978	0.99809	0.98458	0.89758	0.59553
5	0.99998	0.99981	0.99715	0.96598	0.75770
6	0.99999	0.99998	0.99954	0.98989	0.87203

For all $n \geq 2$ we have

$$\lim_{\lambda \rightarrow 0} \sum_{i=1}^n e^{-\lambda \cdot 1} \lambda^{1 \cdot (i-1)} / ((i-1)!)^1 = 1 \text{ and } \lim_{\lambda \rightarrow \infty} \sum_{i=1}^n e^{-\lambda d} \lambda^{d(i-1)} / ((i-1)!)^d = 0 \quad \forall d > 0,$$

hence the dimension attains here all values in $(0, 1)$ for $\lambda \in (0, \infty)$ by continuity.

Let \mathbf{p} now be the zeta distribution with $s > 0$. We have $\dim_H \Pi_{\mathbf{p}}(n) = d$, where d is the solution of

$$\zeta(s)^{-d} \sum_{i=1}^n i^{-sd} = 1.$$

We list the first digits of d for some n and s :

n/s	1.5	2	3	4	5
2	0.48999	0.66938	0.85250	0.92844	0.96292
3	0.64468	0.80840	0.93681	0.97675	0.99085
4	0.72165	0.86713	0.96462	0.98947	0.99667
5	0.76813	0.89903	0.97731	0.99433	0.99850
6	0.79946	0.91890	0.98418	0.99659	0.99923

For all $n \geq 2$ we have

$$\lim_{s \rightarrow 1} \zeta(s)^{-d} \sum_{i=1}^n i^{-sd} = 0 \quad \forall d \quad \text{and} \quad \lim_{s \rightarrow \infty} \zeta(s)^{-1} \sum_{i=1}^n i^{-s \cdot 1} = 1,$$

hence the dimension attains here all values in $(0, 1)$ for $s \in (1, \infty)$ as well.

3 Frequency of digits

Let $\mathbf{f}_{\mathbf{p}}(x, n)$ be the frequency of the digit $n \in \mathbb{N}$ in the representation of $x \in [0, 1)$, given by a probability distribution \mathbf{p} , this means

$$\mathbf{f}_{\mathbf{p}}(x, n) = \lim_{i \rightarrow \infty} \text{Card}\{j | \pi_{\mathbf{p}}^{-1}(x)_j = n \mid j = 1, \dots, i\} / i,$$

if the limit exists. As expected we have

Theorem 3.1 *Let \mathbf{p} be a probability distribution supported by \mathbb{N} . For almost all $x \in [0, 1)$ and all $n \in \mathbb{N}$ we have $\mathbf{f}_{\mathbf{p}}(x, n) = p_n$.*

Proof. Let $\sigma : [0, 1) \rightarrow [0, 1)$ be the piecewise linear expanding map, given by T_n^{-1} on $T_n([0, 1))$ for $n \in \mathbb{N}$. The measure \mathbf{p} on \mathbb{N} induces a Bernoulli measure b on $\mathbb{N}^{\mathbb{N}}$. It is well known (and easy to prove) that this measure is ergodic with respect to the shift map $s(n_k) = n_{k+1}$ on $\mathbb{N}^{\mathbb{N}}$. We refer here to [5] or [9] for introduction to ergodic theory. The map $\pi_{\mathbf{p}}$ projects b to the Lebesgue measure ℓ on $[0, 1)$, $\ell = b \circ \pi_{\mathbf{p}}^{-1}$. Since b is ergodic with respect to s and $\sigma \circ \pi_{\mathbf{p}} = \pi_{\mathbf{p}} \circ s$, the Lebesgue measure ℓ is ergodic with respect

to σ . Applying Birkoff's ergodic theorem to characteristic function χ_n of the interval $T_n([0, 1)) = [\hat{p}_n, p_n + \hat{p}_n)$, we obtain

$$\lim_{i \rightarrow \infty} \frac{1}{i} \sum_{j=1}^i \chi_n(\sigma^j(x)) = p_n$$

for almost all $x \in [0, 1)$ with respect to ℓ . We have $\chi_n(\sigma^j(x)) = 1$ if and only if $\pi_{\mathbf{p}}^{-1}(x)_j = n$. Hence $\mathbf{f}_{\mathbf{p}}(x, n) = p_n$. \square

This theorem has the following immediate corollary, which reminds us on the classical theory of continued fractions:

Corollary 3.1 *For almost all x the representation of x with respect to \mathbf{p} is unbounded.*

Now we consider subset of $[0, 1)$ with prescribed frequencies of digits with respect to a representation given by \mathbf{p} . Let $\mathbf{q} = (q_i)_{i \in \mathbb{N}}$ be a probability distribution on \mathbb{N} , not necessary supported by \mathbb{N} , this means $q_i \in [0, 1]$. We define sets with frequencies of digits given by \mathbf{q} in the following way

$$F(\mathbf{p}, \mathbf{q}) = \{x \in [0, 1) \mid \mathbf{f}_{\mathbf{p}}(x, n) = q_n \ \forall n \in \mathbb{N}\}.$$

Recall that the entropy of \mathbf{q} is

$$H(\mathbf{q}) = - \sum_{i=1}^{\infty} q_i \log(q_i).$$

provided that the limit exists. Here we set $q_i \log(q_i) = 0$ if $q_i = 0$. See [9] or [5] for an introduction to entropy theory. Moreover let

$$E(I_{\mathbf{p}}(\mathbf{q})) = - \sum_{i=1}^{\infty} q_i \log(p_i)$$

provided that the limit exists. This is the expectation of the information content of \mathbf{p} with respect to \mathbf{p} . With these notations we have

Theorem 3.2 *For all probability distributions \mathbf{p} and \mathbf{q} on \mathbb{N} , where the first distribution is supported by \mathbb{N} , we have*

$$\dim_H F(\mathbf{p}, \mathbf{q}) = H(\mathbf{q})/E(I_{\mathbf{p}}(\mathbf{q})),$$

provided that $H(\mathbf{q})$ and $E(I_{\mathbf{p}}(\mathbf{q}))$ exists.

Proof. Let b be the Bernoulli measure, given by \mathbf{q} on $\mathbb{N}^{\mathbb{N}}$. Project this measure to $[0, 1)$, using $\pi_{\mathbf{p}}$, $\mu = b \circ \pi_{\mathbf{p}}^{-1}$. Note that by the law of large numbers we have $\mu(F(\mathbf{p}, \mathbf{q})) = 1$.

For $x \in F(\mathbf{p}, \mathbf{q})$ let $I_{n_1 n_2 \dots n_k}(x)$ be the interval of the form $T_{n_1} \circ T_{n_2} \circ \dots \circ T_{n_k}([0, 1])$ that contains x . By the definition of $F(\mathbf{p}, \mathbf{q})$ we have

$$\begin{aligned} & \lim_{k \rightarrow \infty} \frac{1}{k} \log \frac{\mu(I_{n_1 \dots n_k}(x))}{\text{Length}(I_{n_1 \dots n_k}(x))^s} \\ &= \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{i=1}^k \log(q_{n_i}) - s \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{i=1}^k \log(p_{n_i}) \\ &= -H(\mathbf{q}) + sE(I_{\mathbf{p}}(\mathbf{q})) \end{aligned}$$

provided that $H(\mathbf{q})$ and $E(I_{\mathbf{p}}(\mathbf{q}))$ exist. This implies that for all $x \in F(\mathbf{p}, \mathbf{q})$

$$\lim_{k \rightarrow \infty} \frac{\mu(I_{n_1 \dots n_k}(x))}{\text{Length}(I_{n_1 \dots n_k}(x))^s} = \begin{cases} 0 & s < d \\ \infty & s > d \end{cases},$$

where

$$d = H(\mathbf{q})/E(I_{\mathbf{p}}(\mathbf{q})).$$

By the local mass distribution principle, see proposition 4.9 of [1], we have $\mathfrak{H}^s(F(\mathbf{p}, \mathbf{q})) = \infty$ for $s < d$ and $\mathfrak{H}^s(F(\mathbf{p}, \mathbf{q})) = 0$ for $s > d$. This implies $\dim_H(F(\mathbf{p}, \mathbf{q})) = d$. \square

As an example we consider the set of numbers $F(\mathbf{p}, \mathbf{q})$, which have equidistribution digits from $\{1, \dots, n\}$ in their representation, given by \mathbf{p} . In this case \mathbf{q} is given by $q_i = 1/n$ for $i = 1, \dots, n$. Hence $H(\mathbf{q}) = \log(n)$ and

$$\dim_H F(\mathbf{p}, \mathbf{q}) = -n \log(n) / \log(p_1 \cdots p_n).$$

For the geometric distribution \mathbf{p} , this gives

$$\dim_H F(\mathbf{p}, \mathbf{q}) = \log(n) / ((1 - n) \log(p) - \log(1 - p)),$$

where $p \in (0, 1)$. For the Poisson distribution \mathbf{p} with $\lambda > 1$ we have

$$\dim_H F(\mathbf{p}, \mathbf{q}) = \log(n) / (\lambda - (n - 1) \log(\lambda) / 2 + \sum_{i=1}^{n-1} (n - i) \log(i))$$

and for the zeta distribution \mathbf{p} with $s > 1$ we obtain

$$\dim_H F(\mathbf{p}, \mathbf{q}) = \log(n) / (\log(\zeta(s)) + s \sum_{i=1}^n \log(i) / n).$$

References

- [1] K. Falconer, *Fractal Geometry - Mathematical Foundations and Applications*, Wiley, New York, 1990.
- [2] H. Fernau, *Infinite iterated function systems*, Math. Nach., vol. 170, issue 1, 79-91, 1994.
- [3] J.E. Hutchinson, *Fractals and self-similarity*, Indiana Univ. Math. J. 30, 271-280, 1981.
- [4] V. Jarnik, *Zur metrischen Theorie der diophantischen Approximationen*, Prace Matematyczno-Fizyczne 36.1, 91-106, 1928/29.
- [5] A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems*, Cambridge University Press, 1995.
- [6] P. Moran, *Additive functions of intervals and Hausdorff measure*, Proc. Camb. Phil. Soc. 42, 15-23, 1946.
- [7] R.D. Mauldin and M. Urbanski, *Dimensions and measures in infinite iterated function systems*, Proc. London Math. Soc., 73, 105-154, 1996.
- [8] Ya. Pesin, *Dimension theory in dynamical systems: contemporary view and applications*, The University of Chicago Press 1997.
- [9] P. Walters, *An introduction to ergodic theory*, Springer-Verlag, New York, 1982