

# Non-uniform expansions of real numbers

Jörg Neunhäuserer  
joerg.neunhaeuserer@web.de

## Abstract

We introduce and study non-uniform expansions of real numbers, given by two non-integer bases.

**MSC 2010:** 11K55, 37A45, 28A78

**Key-words:** expansions of real numbers, non-inter bases, cardinality of expansions, unique expansions

## 1 Introduction

Expansions of real numbers in non-integer bases are studied since the pioneering works of Renyi in the end of the 1950s and Parry in the 1960th, see [11, 12, 13]. In these works especially the greedy algorithm that determines the digits of such expansions and the relationship of these expansions to symbolic dynamics is addressed. In the 1990s a group of Hungarian mathematicians led by Paul Erdős revived this field of research, see [2, 4, 5]. Beside other results they proved that each  $x \in (0, 1/(1 - q))$  has a continuum of expansions of the form  $\sum_{i=1}^{\infty} q^{-n_i}$  if  $1 < q < G$ , where  $G$  is the golden ratio. In the sequel Sidorov [14] used ergodic theoretical methods to prove that for all  $q \in (1, 2)$  almost all  $x \in (0, 1/(1 - q))$  have such an expansion. Moreover Glendinning and Sidorov [7] proved that there always exist (at least countably many) reals having a unique expansion if  $q > G$ . Nowadays especially dimensional theoretical aspects of expansions of real numbers in non-integer bases are studied, see for instance [9, 10, 1].

In this paper we introduce non-uniform expansions of real numbers, which may be viewed as expansions with respect to two non-integer bases. As far as we know such expansions were not studied yet, although they constitute a natural generalisation. The rest of the paper is organized as follows:

In the next section we give two descriptions of non-uniform expansions of real numbers. In the following we introduce a greedy, a lazy and intermitted algorithms that give the digits of these expansions. In section four we prove a theorem on the existence of a continuum of non-uniform expansions of real numbers, which is similar to the results in the uniform case we mentioned above. In the last section we characterise real numbers which have a unique non-uniform expansion and prove a theorem on the cardinality of the set of such numbers.

## 2 The expansions

Let  $\Sigma = \{0, 1\}^{\mathbb{N}}$  be the set of sequences of zeros and ones. Equipped with the metric

$$d((s_i), (t_i)) = \sum_{i=1}^{\infty} |s_i - t_i| 2^{-i}$$

$\Sigma$  is a compact, perfect and totally disconnected space. For  $s = (s_i) \in \Sigma$  and  $n \in \mathbb{N}$  let  $0_n(s)$  the number of zeros and  $1_n(s)$  be the number of ones in the sequence  $(s_1, \dots, s_n)$ .

Fix  $\beta_0, \beta_1 \in (1/2, 1)$  with  $\beta_0 \geq \beta_1$  and let  $I = I_{\beta_1} = [0, \beta_1/(1 - \beta_1)]$ .<sup>1</sup> We consider the map  $\pi_{\beta_0, \beta_1} : \Sigma \rightarrow I$  given by

$$\pi_{\beta_0, \beta_1}(s) = \pi_{\beta_0, \beta_1}((s_i)) = \sum_{i=1}^{\infty} s_i \beta_0^{0_n(s)} \beta_1^{1_n(s)}.$$

Several times we will use another description of this map, which we now describe. Let  $T_0, T_1 : I \rightarrow I$  be the contractions given by

$$T_0(x) = \beta_0 x \quad \text{and} \quad T_1(x) = \beta_1 x + \beta_1.$$

By induction we have

$$T_{s_1} \circ \dots \circ T_{s_n}(x) = \beta_0^{0_n(s)} \beta_1^{1_n(s)} x + \sum_{i=1}^n s_i \beta_0^{0_n(s)} \beta_1^{1_n(s)}.$$

Hence for all  $s \in \Sigma$  and all  $x \in I$

$$\pi_{\beta_0, \beta_1}(s) = \pi_{\beta_0, \beta_1}((s_i)) = \lim_{n \rightarrow \infty} T_{s_1} \circ \dots \circ T_{s_n}(x).$$

**Definition 2.1** We call a sequence  $s \in \Sigma$  with  $\pi_{\beta_0, \beta_1}(s) = x$  a  $(\beta_0, \beta_1)$ -expansion of  $x \in I$ .

The following proposition guarantees the existence of  $(\beta_0, \beta_1)$ -expansions.

**Proposition 2.1** *The map  $\pi_{\beta_0, \beta_1}$  is continuous and surjective.*

**Proof.** If  $d((s_i), (t_i)) < 2^{-u}$  we have  $s_i = t_i$  for  $i = 1, \dots, u$ , which implies

$$|\pi_{\beta_0, \beta_1}(s_i) - \pi_{\beta_0, \beta_1}(t_i)| < \beta_0^i \beta_1 / (1 - \beta_1).$$

Hence  $\pi_{\beta_0, \beta_1}$  is continuous. Note that

$$T_0(I) \cup T_1(I) = [0, \beta_0 \beta_1 / (1 - \beta_1)] \cup [\beta_1, \beta_1 / (1 - \beta_1)] = I$$

---

<sup>1</sup>In the literature the uniform case  $\beta_1 = \beta_2$  has been studied. Usually the reciprocal of  $\beta_1$  resp.  $\beta_2$  is denoted by  $\beta$ .

since  $\beta_0 + \beta_1 \geq 1$ . This implies

$$\bigcup_{(s_i) \in \{0,1\}^n} T_{s_1} \circ \cdots \circ T_{s_n}(I) = I.$$

Hence for each  $x \in I$  there is sequence  $s \in \Sigma$  such that

$$x \in T_{s_1} \circ \cdots \circ T_{s_n}(I),$$

but this implies  $\pi_{\beta_0, \beta_1}(s) = x$  and  $\pi_{\beta_0, \beta_1}$  is surjective □

In the next section we describe an algorithm which determines one  $(\beta_0, \beta_1)$ -expansion of  $x \in I$ .

### 3 The greedy and the lazy algorithm

Using the notations of last section we define a map  $G : I \rightarrow I$

$$G(x) = \begin{cases} T_0^{-1}(x), & x \notin T_1(I) \\ T_1^{-1}(x), & x \in T_1(I) \end{cases}$$

$$= \begin{cases} \beta_0^{-1}x, & x \in [0, \beta_0) \\ \beta_1^{-1}x - 1, & x \in [\beta_0, \beta_1/(1 - \beta_1)], \end{cases}.$$

For  $x \in I$  we define the greedy expansion  $g = g(x) = (g_i) \in \Sigma$  with respect to  $(\beta_1, \beta_2)$  by

$$g_i = \lfloor \beta_0^{-1} G^{i-1}(x) \rfloor,$$

where  $\lfloor a \rfloor$  is the greatest integer not greater than  $a$ . We have

**Proposition 3.1** *For all  $x \in I$  the greedy expansion  $g(x) \in \Sigma$  with respect to  $(\beta_1, \beta_2)$  is an  $(\beta_0, \beta_1)$ -expansion of  $x$ , that is  $\pi_{\beta_0, \beta_1}(g(x)) = x$ .*

**Proof.** If  $g_i = 0$  we have  $G^{i-1}(x) < \beta_0$ , which implies  $G^{i-1}(x) \notin T_1(I)$  and  $G^{i-1}(x) \in T_{g_i}(I)$ . If  $g_i = 1$  we have  $G^{i-1}(x) \geq \beta_0$ , which implies  $G^{i-1}(x) \in T_1(I)$  hence again  $G^{i-1}(x) \in T_{g_i}(I)$ . By the definition of  $G$  we conclude

$$x \in T_{g_1} \circ \cdots \circ T_{g_i}(I)$$

for all  $i \in \mathbb{N}$ , but this implies  $\pi_{\beta_0, \beta_1}(g(x)) = x$ . □

To define the lazy expansion let  $L : I \rightarrow I$  be given by

$$L(x) = \begin{cases} T_0^{-1}(x), & x \in T_0(I) \\ T_1^{-1}(x), & x \notin T_0(I) \end{cases}$$




Figure 1: The maps  $T_0^{-1}$  and  $T_1^{-1}$  in the case  $\beta_0 = 3/4, \beta_1 = 2/3$

$$= \begin{cases} \beta_0^{-1}x, & x \in [0, \beta_0\beta_1/(1-\beta_1)] \\ \beta_1^{-1}x - 1, & x \in (\beta_0\beta_1/(1-\beta_1), \beta_1/(1-\beta_1)], \end{cases} .$$

For  $x \in I$  the lazy expansion  $l = l(x) = (l_i) \in \Sigma$  with respect to  $(\beta_1, \beta_2)$  is given by

$$l_i = \lceil (1 - \beta_1)(\beta_0\beta_1)^{-1}L^{i-1}(x) \rceil,$$

where  $\lceil a \rceil$  is the smallest integer not smaller than  $a$ . We have

**Proposition 3.2** *For all  $x \in I$  the lazy expansion  $l(x) \in \Sigma$  with respect to  $(\beta_1, \beta_2)$  is an  $(\beta_0, \beta_1)$ -expansion of  $x$ , that is  $\pi_{\beta_0, \beta_1}(l(x)) = x$ .*

**Proof.** If  $l_i = 0$  we have  $L^{i-1}(x) \leq \beta_0\beta_1/(1-\beta_1)$ , which implies  $L^{i-1}(x) \in T_0(I)$  hence  $L^{i-1}(x) \in T_{l_i}(I)$ . If  $l_i = 1$  we have  $L^{i-1}(x) > \beta_0\beta_1/(1-\beta_1)$ , which implies  $L^{i-1}(x) \notin T_0(I)$  hence again  $L^{i-1}(x) \in T_{l_i}(I)$ . By the definition of  $L$  we conclude

$$x \in T_{l_1} \circ \dots \circ T_{l_i}(I)$$

for all  $i \in \mathbb{N}$ , which implies  $\pi_{\beta_0, \beta_1}(l(x)) = x$ . □

For  $\alpha \in (\beta_1, \beta_0\beta_1/(1-\beta_1))$  we may also consider intermediate expansions  $m(x) = (m_i)$  with respect to  $(\beta_1, \beta_2)$  given by

$$m_i = \lfloor \alpha^{-1}M_\alpha^{i-1}(x) \rfloor,$$

where

$$M_\alpha(x) = \begin{cases} \beta_0^{-1}x, & x \in [0, \alpha) \\ \beta_1^{-1}x - 1, & x \in [\alpha, \beta_1/(1 - \beta_1)] \end{cases}.$$

Again these are  $(\beta_0, \beta_1)$ -expansion of  $x \in I$ .

## 4 A continuum of expansions

It is natural to ask how many  $(\beta_0, \beta_1)$ -expansions a real number in  $I$  has. It turns out that usually there is a continuum of such expansions:

**Theorem 4.1** *Let  $\beta_0, \beta_1 \in (1/2, 1)$  with  $\beta_0 \geq \beta_1$ . We have:*

(1) *Almost all  $x \in I$  have a continuum of  $(\beta_0, \beta_1)$ -expansions.*

(2) *If  $\beta_1^2 + \beta_0 > 1$  all  $x \in I \setminus \{0, \beta_1/(1 - \beta_1)\}$  have a continuum of  $(\beta_0, \beta_1)$ -expansions.*

**Proof.** We first prove (2). Let  $J = (0, \beta_1/(1 - \beta_1))$  and

$$\Lambda_0 = T_0(J) \cap T_1(J) = (\beta_1, \beta_0 \frac{\beta_1}{1 - \beta_1}).$$

We recursively define  $\Lambda_{n+1} = T_0(\Lambda_n) \cup \Lambda \cup T_1(\Lambda_n)$  and prove by induction:

$$\Lambda_n = (\beta_0^n \beta_1, (\beta_1^n (\beta_0 - 1) + 1) \frac{\beta_1}{1 - \beta_1}).$$

We have

$$\begin{aligned} T_0(\Lambda_n) \cup \Lambda \cup T_1(\Lambda_n) &= (\beta_0^{n+1} \beta_1, (\beta_1^n \beta_0 (\beta_0 - 1) + 1) \frac{\beta_1}{1 - \beta_1}) \cup (\beta_0^n \beta_1, (\beta_1^n (\beta_0 - 1) + 1) \frac{\beta_1}{1 - \beta_1}) \\ &\cup (\beta_0^n \beta_1^2 + \beta_1, (\beta_1^{n+1} (\beta_0 - 1) + 1) \frac{\beta_1}{1 - \beta_1}) = \Lambda_{n+1}. \end{aligned}$$

In the last equation we use  $\beta_1^2 + \beta_0 > 1$  and  $\beta_0^2 + \beta_1 > 1$ , which is true since  $\beta_0 \geq \beta_1$ . Note that  $\bigcup_{n \geq 0} \Lambda_n = J$ . Hence for every  $x \in J$  there is a  $k \geq 0$  and a sequence  $(s_1, \dots, s_k) \in \{0, 1\}^k$  such that

$$x = T_{s_1} \circ \dots \circ T_{s_k} \circ T_0(x_0) \text{ and } x = T_{s_1} \circ \dots \circ T_{s_k} \circ T_1(x_1),$$

where  $x_0, x_1 \in J$  and  $x_0 \neq x_1$ . Hence we obtain two expansions of  $x$  that differ in the  $k + 1$ -digit. Applying the result to  $x_0(x)$  and  $x_1(x)$  we obtain four expansions of  $x$ . Here we use that  $x_0(x)$  and  $x_1(x)$  are not at the boundary of  $J$ . Repeating this procedure  $\aleph_0$  times we see that there are  $2^{\aleph_0}$  expansions of  $x$ .

Now we prove (1). Let  $G : I \rightarrow I$  be the map associated with the greedy expansion from section 3.  $G$  is a piecewise linear expanding interval map and such maps are known to

have an ergodic measure, which is equivalent to the Lebesgue measure, see [3] and [8]. By Poincare recurrence theorem for almost all  $x \in I$  there is a  $k \geq 0$  such that  $G^k(x) \in \Lambda_0$ . Hence for almost all  $x \in J$  there is a  $k \geq 0$  and a sequence  $(s_1, \dots, s_k) \in \{0, 1\}^k$  such that

$$x = T_{s_1} \circ \dots \circ T_{s_k} \circ T_0(x_0) \text{ and } x = T_{s_1} \circ \dots \circ T_{s_k} \circ T_1(x_1),$$

where  $x_0, x_1 \in J$  and  $x_0 \neq x_1$ . For almost all  $x$  both numbers  $x_1(x), x_2(x)$  have two different  $(\beta_0, \beta_1)$ -expansion hence almost all  $x$  have four different expansions. We use here that the intersection of two sets of full measure has full measure. Repeating this procedure  $\aleph_0$  times we obtain  $2^{\aleph_0}$  expansions for almost all  $x \in I$ , using the fact a countable intersection of sets of full measure has full measure.  $\square$

Obviously the  $(\beta_0, \beta_1)$ -expansion of 0 and  $\beta_1/(1 - \beta_1)$  is unique. Our theorem leaves the question open if there are numbers  $x$  in the interior of  $I$  that have a unique  $(\beta_0, \beta_1)$ -expansion. We will address this question in the following section.

## 5 Unique expansions

We consider the shift map  $\sigma : \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}^{\mathbb{N}}$  given by  $\sigma((s_k)) = (s_{k+1})$ . Using this map we may characterise numbers which have a unique  $(\beta_0, \beta_1)$ -expansion as follows:

**Proposition 5.1** *The  $(\beta_0, \beta_1)$ -expansion  $(s_i)$  of  $x$  is unique if and only if*

$$\pi_{\beta_0, \beta_1}(\sigma^k(s_i)) \in [0, \beta_1] \cup (\beta_0\beta_1/(1 - \beta_1), \beta_1/(1 - \beta_1)]$$

for all  $k \geq 0$ .

**Proof.**  $\pi_{\beta_0, \beta_1}((s_i)) = \pi_{\beta_0, \beta_1}((t_i))$  with  $(s_i) \neq (t_i)$  if and only if there exists a  $k \geq 0$  such that  $s_k \neq t_k$  and  $\pi_{\beta_0, \beta_1}(\sigma^k(s_i)) = \pi_{\beta_0, \beta_1}(\sigma^k(t_i))$ . But this is equivalent to  $\pi_{\beta_0, \beta_1}(\sigma^k(s_i)) \in T_0(I) \cap T_1(I) = [\beta_1, \beta_0\beta_1/(1 - \beta_1)]$ . The proposition follows by contraposition.  $\square$

Using this characterisation of points with unique expansion we are able to prove:

**Theorem 5.1** *Let  $\beta_0, \beta_1 \in (1/2, 1)$  and  $\beta_0 \geq \beta_1$ .*

(1) *If  $\beta_0(1 + \beta_1) < 1$  there are at least countable many  $x \in I$ , which have a unique  $(\beta_0, \beta_1)$ -expansion.*

(2) *If  $\beta_0(1 + 2\beta_1 - \beta_0\beta_1) < 1$  there are uncountable many  $x \in I$ , which have a unique  $(\beta_0, \beta_1)$ -expansion. Moreover the set of these  $x$  has positive Hausdorff dimension.*

**Proof.** Consider the periodic sequence  $p = (010101\dots)$ . Since  $\beta_0(1 + \beta_1) < 1$  we have

$$\pi_{\beta_0, \beta_1}(p) = \beta_0\beta_1/(1 - \beta_0\beta_1) < \beta_1.$$

Note that  $\beta_0(1 + \beta_1) < 1$  implies  $\beta_1(1 + \beta_0) < 1$  since  $\beta_0 \geq \beta_1$ . Hence we have  $\beta_0 - \beta_0^2\beta_1 < 1 - \beta_1$  and thus

$$\pi_{\beta_0, \beta_1}(\sigma(p)) = \beta_1 / (1 - \beta_0\beta_1) = \beta_0\beta_1 / (\beta_0 - \beta_0^2\beta_1) > \beta_0\beta_1 / (1 - \beta_1).$$

By proposition  $x = \pi_{\beta_0, \beta_1}(p)$  has a unique  $(\beta_0, \beta_1)$ -expansion. Obviously the same is true for all  $x$  of the form  $x = \pi_{\beta_0, \beta_1}((0 \dots 0101010 \dots))$  and there are countably many of such  $x$ . Now we prove (2). Let  $V = \{01, 10\}^{\mathbb{N}}$  and

$$U = \bigcup_{k=0}^{\infty} \sigma^k(V) = V \cup (\{0\} \times V) \cup (\{1\} \times V).$$

We prove that  $\pi_{\beta_0, \beta_1}(U) \subseteq [0, \beta_1] \cup (\beta_0\beta_1 / (1 - \beta_1), \beta_1 / (1 - \beta_1)]$ . The sequence  $s \in U$  with  $s_1 = 0$  that has the largest projection under  $\pi_{\beta_0, \beta_1}$  obviously is  $s = (011010101 \dots)$ . We have

$$\pi_{\beta_0, \beta_1}(s) = \beta_1 \frac{\beta_0 + \beta_0\beta_1 - \beta_0^2\beta_1}{1 - \beta_0\beta_1} < \beta_1$$

by our assumption. The sequence  $s \in U$  with  $s_1 = 1$  that has the smallest projection under  $\pi_{\beta_0, \beta_1}$  obviously is  $s = (1001010101 \dots)$ . We have

$$\pi_{\beta_0, \beta_1}(s) = \beta_1 + \frac{(\beta_0\beta_1)^2}{1 - \beta_0\beta_1} > \frac{\beta_0\beta_1}{1 - \beta_1}.$$

The inequality here is equivalent to  $\beta_1(1 + 2\beta_0 - \beta_0\beta_1) < 1$  which is true since we assume  $\beta_0 \geq \beta_1$ . It remains to show that the Hausdorff dimension of  $A := \pi_{\beta_0, \beta_1}(V)$  is positive. Consider the maps

$$F(x) = T_0 \circ T_1(x) = \beta_0\beta_1x + \beta_0\beta_1$$

and

$$G(x) = T_1 \circ T_0(x) = \beta_0\beta_1x + \beta_1$$

and let  $J = (0, \beta_1 / (1 - \beta_0\beta_1))$ . We have  $F(J) \subseteq J$  and  $G(J) \subseteq J$ . Moreover

$$F(J) \cap G(J) = (0, \beta_0\beta_1^2 / (1 - \beta_0\beta_1)) \cap (\beta_1, \beta_1 / (1 - \beta_0\beta_1)) = \emptyset$$

by our assumptions on  $\beta_0$  and  $\beta_1$ . In the language of fractal geometry this means that  $(F, G)$  induce an iterated function system fulfilling the open set condition, see [6]. The attractor of this iterated function system is  $A$  since  $A = F(A) \cup G(A)$  and the classical formula for self-similar fractals gives

$$\dim_H A = \frac{-\log(2)}{\log(\beta_0\beta_1)} > 0.$$

□

## References

- [1] R. Alcaraz Barrera, S. Baker, Simoe and D. Kong, Entropy, topological transitivity, and dimensional properties of unique  $q$ -expansions, *Trans. Am. Math. Soc.* 371 (2019), No. 5, 3209-3258.
- [2] P. Erdős, I. Joo and V. Komornik, Characterization of the unique expansions  $\sum_{i=1}^{\infty} q^{-n_i}$  and related problems, *Bull. Soc. Math. Fr.* 118 (1990), 377-390.
- [3] M. Denker, *Introduction to analysis of dynamical systems*, Springer, Berlin, 2005.
- [4] P. Erdős, I. Joo, On the number of expansions  $\sum_{i=1}^{\infty} q^{-n_i}$ , *Ann. Univ. Sci. Budapest* 35 (1992), 129-132.
- [5] P. Erdős, I. Joo and V. Komornik, On the number of  $q$ -expansions, *Ann. Univ. Sci. Budapest Eötvös Sect. Math.* 37 (1994), 109-118.
- [6] K. Falconer, *Fractal Geometry - Mathematical Foundations and Applications*, Wiley, New York, 1990.
- [7] P. Glendinning and N. Sidorov, Unique representations of real numbers in non-integer bases, *Math. Res. Letters* 8 (2001), 535-543.
- [8] A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems*, Cambridge University Press, 1995.
- [9] V. Komornik, D. Kong and W., Hausdorff dimension of univoque sets and devils staircase, *Adv. Math.* 305 (2017), 165-196.
- [10] D. Kong, W. Li; F. Lü and M. de Vries, Univoque bases and Hausdorff dimension. *Monatsh. Math.* 184 (2017), No. 3, 443-458.
- [11] A. Renyi, Representations for real numbers and their ergodic properties, *Acta Math. Acad. Sci. Hung.* 8 (1957) 477-493.
- [12] W. Parry, On the  $\beta$ -expansions of real numbers, *Acta Math. Acad. Sci. Hung.* 11 (1960) 401-416.
- [13] W. Parry, Representations for real numbers, *Acta Math. Acad. Sci. Hungar.* 15 (1964), 95-105.
- [14] N. Sidorov, Almost every number has a continuum of  $\beta$ -expansions, *Amer. Math. Monthly* 110 (2003), 838-842.